

The potential of INULIN as a PREBIOTIC in pets



PROMOTES GASTROINTESTINAL HEALTH

- Positively affects indices associated with gut health¹
- Boosts concentrations of faecal short-chain fatty acids^{1,2,3}
- Promotes laxation⁴
- Reduces level of ammonia (likely improving stool odour)⁵

HELPS DIGESTION

- Positively affects apparent protein digestibility^{6,7}



IMPROVES METABOLISM

- Improves blood glucose and lipid metabolism in dogs with obesity⁸
- Indirectly stimulates whole-body acetate turnover⁹

POTENTIAL IN AMELIORATING OBESITY

- Mitigates weight gain and helps maintain body condition score in overfed dogs⁸



REGULATES MICROBIOTA

- Reduces abundance of potentially harmful *Enterobacteriaceae* and *Escherichia coli*⁸
- Increases beneficial bacteria such as *Lactobacillus* and *Akkermansia*¹⁰

REDUCES GASTROINTESTINAL INFLAMMATION

- Inhibits a key inflammatory signaling pathway & inflammatory factors in obese dogs⁸



To find out more, go to Purina.co.uk/Inulin






PURINA[®] PRO PLAN[®]

REFERENCES

1. Propst, E. L., Flickinger, E. A., Bauer, L. L., Merchen, N. R., & Fahey Jr, G. C. (2003). A dose-response experiment evaluating the effects of oligofructose and inulin on nutrient digestibility, stool quality, and fecal protein catabolites in healthy adult dogs. *Journal of animal science*, 81 (12), 3057-3066.
2. Alexander, C., Cross, T. W. L., Devendran, S., Neumer, F., Theis, S., Ridlon, J. M., ... & Swanson, K. S. (2018). Effects of prebiotic inulin-type fructans on blood metabolite and hormone concentrations and faecal microbiota and metabolites in overweight dogs. *British Journal of Nutrition*, 120(6), 711-720
3. Hesta, M., Janssens, G. P. J., Debraekeleer, J., & De Wilde, R. (2001). The effect of oligofructose and inulin on faecal characteristics and nutrient digestibility in healthy cats. *Journal of Animal Physiology and Animal Nutrition*, 85(5-6), 135-141.
4. Strompfová, V., Lauková, A., & Cílik, D. (2013). Synbiotic administration of canine-derived strain *Lactobacillus fermentum* CCM 7421 and inulin to healthy dogs. *Canadian journal of microbiology*, 59(5), 347-352.
5. Vierbaum, L., Eisenhauer, L., Vahjen, W., & Zentek, J. (2019). In vitro evaluation of the effects of *Yucca schidigera* and inulin on the fermentation potential of the faecal microbiota of dogs fed diets with low or high protein concentrations. *Archives of animal nutrition*, 73(5), 399-413.
6. Verlinden, A., Hesta, M., Hermans, J. M., & Janssens, G. P. J. (2006). The effects of inulin supplementation of diets with or without hydrolysed protein sources on digestibility, faecal characteristics, haematology and immunoglobulins in dogs. *British Journal of Nutrition*, 96(5), 936-944.
7. Barry, K. A., Hernot, D. C., Middelbos, I. S., Francis, C., Dunsford, B., Swanson, K. S., & Fahey Jr, G. C. (2009). Low-level fructan supplementation of dogs enhances nutrient digestion and modifies stool metabolite concentrations, but does not alter fecal microbiota populations. *Journal of animal science*, 87(10), 3244-3252.
8. Lu, J., Zhu, D., Lu, J., Liu, J., Wu, Z., & Liu, L. (2022). Dietary supplementation with low and high polymerization inulin ameliorates adipose tissue inflammation via the TLR4/NF- κ B pathway mediated by gut microbiota disturbance in obese dogs. *Research in Veterinary Science*, 152, 624-632.
9. Pouteau, E., Frenais, R., Dumon, H., Noah, L., Martin, L., & Nguyen, P. (2005). Colonic fermentation of inulin increases whole-body acetate turnover in dogs. *The Journal of nutrition*, 135(12), 2845-2851.
10. Beloshapka, A. N., Dowd, S. E., Suchodolski, J. S., Steiner, J. M., Duclos, L., & Swanson, K. S. (2013). Fecal microbial communities of healthy adult dogs fed raw meat-based diets with or without inulin or yeast cell wall extracts as assessed by 454 pyrosequencing. *FEMS microbiology ecology*, 84(3), 532-541.



To find out more, go to Purina.co.uk/Inulin

